

MAGNETIC FLARES ON ASYMPTOTIC GIANT BRANCH STARS

Noam Soker^{1,2} and Joel H. Kastner¹

ABSTRACT

We investigate the consequences of magnetic flares on the surface of asymptotic giant branch (AGB) and similar stars. In contrast to the solar wind, in the winds of AGB stars the gas cooling time is much shorter than the outflow time. As a result, we predict that energetic flaring will not inhibit, and may even enhance, dust formation around AGB stars. If magnetic flares do occur around such stars, we expect some AGB stars to exhibit X-ray emission; indeed certain systems including AGB stars, such as Mira, have been detected in X-rays. However, in these cases, it is difficult to distinguish between potential AGB star X-ray emission and, e.g., X-ray emission from the vicinity of a binary companion. Analysis of an archival ROSAT X-ray spectrum of the Mira system suggests an intrinsic X-ray luminosity $\sim 2 \times 10^{29}$ erg s⁻¹ and temperature $\sim 10^7$ K. These modeling results suggest that magnetic activity, either on the AGB star (Mira A) or on its nearby companion (Mira B), is the source of the X-rays, but do not rule out the possibility that the X-rays are generated by an accretion disk around Mira B.

Subject headings: stars: mass loss — stars: winds, outflows — X-rays: ISM — stars: AGB — stars: magnetic fields

1. INTRODUCTION

Asymptotic giant branch (AGB) stars represent the final stages of evolution for intermediate mass ($\sim 1 - 8M_{\odot}$) stars, just before these stars (potentially) generate planetary nebulae. Such stars have exhausted their core hydrogen, have completed core helium and

¹Chester F. Carlson Center for Imaging Science, Rochester Institute of Technology, 54 Lomb Memorial Dr., Rochester, NY 14623; jhk@cis.rit.edu

²Department of Physics, Oranim, Tivon 36006, ISRAEL; soker@physics.technion.ac.il

H-shell burning phases, and are now burning H and He in concentric shells around the exhausted stellar core (this core, once exposed, will become a degenerate white dwarf). AGB stars are ascending fully convective Hayashi tracks in the H-R diagram, and the very large luminosities of these stars ($\sim 10^4 L_\odot$), in combination with copious dust grain formation in their cool, extended atmospheres, lead to dusty, radiatively-driven winds. Hence, AGB stars lose mass rapidly, with rates ranging from $\sim 10^{-8}$ to $\sim 10^{-4} M_\odot \text{ yr}^{-1}$ (as ascertained from far-IR continuum and millimeter-wave molecular line emission; e.g. Loup et al. 1993).

There are several independent pieces of evidence for the presence of magnetic fields in and around cool giants and, in particular, in and around AGB stars. These include:

1. Maser polarization around AGB stars and other cool stars. SiO masers are found close to (typically within less than a few stellar radii of) the stellar surface (e.g. Kemball & Diamond 1997), OH masers are found at $\sim 10^{15} - 10^{16}$ cm from the star (e.g., Szymczak, Cohen, & Richards 1999; Palen & Fix 2000; Bains et al. 2003), and H₂O masers are found in between (Vlemmings, Diamond, & van Langevelde 2002).
2. Polarization of OH maser spots in planetary nebulae (PNs) and proto-PNs, which are the immediate descendants of AGB stars (e.g., Zijlstra et al. 1989). Miranda et al. (2001) find polarization in the 1,665-MHz OH maser line, which indicates the presence of $\sim 10^{-3}$ G magnetic fields at $\sim 10^{16}$ cm from the central star of the young PN K3-35.
3. X-ray emission from cool giant stars (Hünsch et al. 1998; Schröder, Hünsch, & Schmitt 1998; Hünsch 2001). Hünsch (2001) used *Chandra* to detect X-ray emission from two M giant stars, which display hard X-ray luminosities of $L_x \simeq 10^{30} \text{ erg s}^{-1}$.

Although it is widely agreed that radiation pressure on dust is the main acceleration process of winds from AGB stars (Elitzur, Ivezić, & Vinković 2002, and references therein), there is still disagreement over the role the magnetic field plays in shaping the circumstellar matter. First, it is not clear whether magnetic fields play any role in shaping the winds; many other models exist, e.g., acceleration via radiation pressure on dust in fast rotating AGB stars spun up by companions (e.g., Dorfi & Höfner 1996; Reimers, Dorfi & Höfner 2000), and the influence of a companion outside the envelope (e.g., Mastrodemos & Morris 1999; Soker 2001). Second, there is a disagreement on whether the magnetic fields play a dynamical role, i.e., whether the magnetic pressure and/or tension become comparable to the other relevant forces in the flow (e.g., Pascoli 1997; Chevalier & Luo 1994; García-Segura 1997; García-Segura et al. 1999; García-Segura, & López 2000; García-Segura, López, & Franco 2001; Matt et al. 2000; Blackman et al. 2001; Gardiner & Frank 2001; Falceta-Goncalves & Jatenco-Pereira 2002; see review by García-Segura 2002), or whether the magnetic field

is relatively weak on a global scale, and plays only secondary effects (Soker 2002; Soker & Zoabi 2002 and references therein).

In earlier papers Soker and collaborators (e.g., Soker 1998; Soker & Clayton 1999) propose that cool magnetic spots facilitate the formation of dust, hence locally increasing the mass loss rate. In this process the magnetic field has a secondary role and becomes dynamically important only in relatively small regions of the stellar surface and circumstellar environment. In this model the large scale magnetic field is too weak to play a dynamic role and directly influence the wind from the AGB star. However, the magnetic field is strong enough above and near cool spots that reconnection may occur (if magnetic pressure exceeds thermal pressure); such reconnection events likely would lead to magnetic flares analogous to those on the surface of the sun. The reconnection process is basically cancellation of magnetic field lines having opposite directions. The energy released by the reconnecting magnetic fields heats the gas and accelerates it to high speeds. The gas is heated by MHD shocks (e.g., Chen et al. 1999). In the present paper we discuss some implications of these events. In §2 the potential of impact of flaring on circumstellar dust formation is considered. In §3 we discuss the expected observational signatures of AGB star magnetic activity in general and of flares in particular. Our summary is in §4.

2. DUST FORMATION IN FLARES

Whereas Soker (1998) and Soker & Clayton (1999) argued that magnetic starspots on cool giants can lead to enhanced dust formation, we demonstrate here that reconnection events (flares) associated with such spots may themselves further enhance the formation rate of dust. Because the density close to the surface of AGB stars is very high, the question regarding dust formation reduces to the value of the temperature. To evaluate the temperature resulting from a flare, the time scale of cooling should be compared with the flow time. In the sun the conduction time is non-negligible compared with the radiative cooling time (e.g., Vesecky, Antiochos, & Underwood 1979; Aschwanden et al. 2000). However, in AGB stars it can be neglected. This can be shown by using equations (16) and (17) from Aschwanden et al. 2000: taking a temperature of $T \sim 10^6$ K and electron density of $n_e \sim 10^{10}$ in their equations (16) and (17) yields a radiative cooling time of $\tau_{\text{rad}} \sim 800$ s (see below), and a conduction time scale of $\tau_{\text{cond}} \sim 10^9 (L/10^{12} \text{ cm})^2 \text{ s} \gg \tau_{\text{rad}}$, where L is the loop length.

Let us consider an idealized simple case, in which the magnetic field expands radially from the stellar surface outward, while the density is determined by an isothermal hydrostatic halo. In such a case, the magnetic field decreases relatively slowly, $B(r) = B_*(r/R_*)^{-2}$, where

B_* is the magnetic field on the stellar surface, and R_* is the stellar radius (subscript $*$ will stand for quantities on the stellar surface). For an isothermal gas above and close to the photosphere the density is given by

$$\rho = \rho_* \exp[-(r - R_*)/H_*], \quad (1)$$

where the scale height close to the stellar surface is

$$\frac{H}{R_*} \equiv \frac{C_*^2}{v_K^2} \simeq 0.045 \left(\frac{T_*}{3000 \text{ K}} \right) \left(\frac{R_*}{1 \text{ AU}} \right) \left(\frac{M_*}{M_\odot} \right)^{-1}. \quad (2)$$

In equation (2), C_* and v_K are the isothermal sound speed and Keplerian velocity on the stellar surface, respectively. The temperature was scaled for pre-flare gas because we do not expect AGB stars to possess long lived coronae. In any case, in calculating C_* we took a fully ionized gas so the equation can be used for higher temperatures. For reconnection to occur and heat the gas substantially, the magnetic pressure should overwhelm the thermal pressure. This occurs several scale heights above the surface, and higher. In the case of the sun the same condition is met very close to the solar surface, and we can have small flares as well as large flares. The smaller flares (nanoflares) above the solar surface have minimum height of $h \sim 0.0015R_\odot \simeq 4H_\odot$ (e.g., Aschwanden et al. 2000). Inside magnetic flux loops the magnetic pressure is high, and pressure equilibrium with the loop exterior implies that the density inside the loop is smaller than in its surroundings. Also, after heating following reconnection, the hot gas expands and the density drops below the simple isothermal halo value. Strong flares extend to much larger distances from the solar surface, $h \simeq 0.007 - 0.07R_\odot \simeq 20 - 200H_\odot$ (e.g., Aschwanden et al. 2000).

As can be seen from equation (2), on the surface of AGB stars, for which $R_* \gtrsim 1 \text{ AU}$, the scale height is $H \sim 0.05R_*$. As just noted for the case of the sun, flares, from small to large, extend from $h \sim 4H$ to $h \sim 200H$ above the solar surface. Taking the same range for AGB stars, for which $H \simeq 0.05R_*$ (Eq. 2), we find that flares will occur over the range of $r = R_* + h \simeq 1.2 - 10R_*$. Since in the upper range $h \gg R_*$, it is not clear if flares can extend to such large radii above the surfaces of AGB stars. However, there is no physical reason why flares will not occur on small scales, i.e., $h \sim 0.2 - 1R_*$. The magnetic energy can even be released in several consecutive reconnection events, without changing our conclusions. We also note that even in large flares the total magnetic energy is still much below that required in models where the magnetic field has a dynamical role (see Section 1).

The typical density of the photospheres of AGB stars is $n_e(R_*) \sim 10^{14} \text{ cm}^{-3}$ (Soker & Harpaz 1999³). In the simple isothermal halo used above, the density drops to $n_e \simeq 10^{10} \text{ cm}^{-3}$

³Note that the density scale in Figs. 1-5 of Soker & Harpaz 1999 is too low by a factor of 10; the correct

at $r \sim 1.3R_*$. At distances of $r \gtrsim 2R_*$ we are in the region where density is determined by the wind, i.e., $\rho = \dot{M}/(4\pi r^2 v_w)$, where $\dot{M}$ is the mass loss rate into the wind (defined positively), and $v_w(r)$ is the wind speed. For a fully ionized wind — which is not the case for the global AGB wind, but will be the case locally after a flare event — the scaled electron density is

$$n_e \simeq 10^{10} \left(\frac{\dot{M}}{10^{-6} M_\odot \text{ yr}^{-1}} \right) \left(\frac{r}{1 \text{ AU}} \right)^{-2} \left(\frac{v_w}{10 \text{ km s}^{-1}} \right)^{-1} \text{ cm}^{-3} \quad (3)$$

where, for simplicity, we have assumed isotropic mass loss. Using the cooling function from, e.g., Gaetz, Edgar & Chevalier (1988), we find the isobaric radiative cooling time in the temperature range $2 \times 10^5 \text{ K} \lesssim T \lesssim 4 \times 10^7 \text{ K}$ to be

$$\tau_{\text{rad}} \simeq 800 \left(\frac{n_e}{10^{10}} \right)^{-1} \left(\frac{T}{10^6} \right)^{3/2} \text{ s}. \quad (4)$$

The flow time across the reconnection region can be estimated as follows. Let the size of the reconnection flux tube be $x \simeq H \sim 10^{12} \text{ cm}$, and the flow speed be the sound speed at $T = 10^6 \text{ K}$, i.e., $v_{\text{exp}} \sim 100 \text{ km s}^{-1}$. The expansion time is then

$$\tau_{\text{exp}} \sim 10^5 \left(\frac{x}{H} \right) \left(\frac{H}{10^{12} \text{ cm}} \right) \left(\frac{v_e}{100 \text{ km s}^{-1}} \right)^{-1} \text{ s}. \quad (5)$$

Even if $x \sim 0.1H$, the radiative cooling time of gas heated by flares to $T \sim 10^6 \text{ K}$ in AGB stars is much shorter than the expansion time. The cooling time will also be shorter than the expansion time in strong and large flares, having a temperature of $T \sim 10^7 \text{ K}$ and a size of $x \gtrsim 10H$. Close to the AGB stellar surface the wind does not yet reach its terminal velocity, hence the density is higher and the cooling time shorter than the values in equations (3) and (4), respectively. In the sun, where $H_* \sim 3 \times 10^7 \text{ cm}$, we find that the situation is different, with the radiative cooling time much longer than the expansion time (see, e.g., Aschwanden et al. 2000). At larger distances from the surface of AGB stars, $r \sim 10R_*$, the cooling time is no longer much shorter than the expansion time; nevertheless, the condition that the cooling time is much shorter than the expansion time holds for dust formation close to the stellar surface.

The compression in the strong shock that results from the reconnection event can be larger than a factor of 4. This is because the gas is not fully ionized. For conditions above cool giants, and for a shock temperature of $\sim 10^6 \text{ K}$, the post shock density can be a factor

scale is displayed in their Fig. 6.

$\sim 5 - 10$ larger than the pre-shock density (Woitke, Gores & Sedlmayr 1996a). Since the gas cools much faster than expansion time, to maintain pressure equilibrium the gas will be further compressed. The situation here is more complicated than the shocks generated by pulsating stars, as discussed by Woitke et al. (1996a), because the shocks are MHD shocks (e.g., Chen et al. 1999). After the gas cools from $T \sim 10^6$ to $T \sim 10^4$ K, we can take, therefore, a density that is larger by a factor of ~ 10 relative to the value in equation (3).

After this gas cools to $T \simeq 10^4$ K, the radiative cooling time increases substantially (e.g., Woitke, Krüger, & Sedlmayr 1996b); on the other hand, the cooling now proceeds radiatively as well as via adiabatic expansion (Woitke et al. 1996a). To form dust, the temperature should decrease to $T \lesssim 10^3$ K. Using figure (11) of Woitke et al. (1996b), we approximate the cooling time from a temperature of $T \simeq 10^4$ K down to dust-forming temperatures by

$$\tau_{\text{rad}} \simeq 5 \times 10^6 \left(\frac{n_H}{10^{10} \text{ cm}^{-3}} \right)^{-0.3} \text{ s}, \quad (6)$$

where n_H is the hydrogen number density. Note that the gas is not fully ionized in this case, in contrast to the temperature range for which equation (4) holds. The cooling time is still short, however. The cooler gas will expand at slower velocity; either the sound speed or the wind speed give $v_e \simeq 10 \text{ km s}^{-1}$, which gives for the expansion time $\tau_e \sim 10^6$ s, according to equation (5). Taking $n_H \sim 10^{11} \text{ cm}^{-3}$, as discussed above, and considering further cooling by adiabatic expansion (Woitke et al. 1996a), we find the expansion time of the gas to be comparable to the cooling time for the scaling used above, i.e., a mass loss rate of $\dot{M} = 10^{-6} M_\odot \text{ yr}^{-1}$ in equation (3).

The main conclusion from this discussion is that magnetic flares on the surfaces of AGB stars, as well as on other classes of actively mass-losing cool giant stars which lose mass at a high rate, will not inhibit dust formation, because the gas cools on a short time scale. The basic mechanism is that the lower temperature above cool magnetic spots allows dust to form closer to the AGB surface where the density is higher, thereby enhancing dust formation and, consequently, mass loss rate (Soker 1998, 2000; Soker & Clayton 1999). Furthermore, strong shocks, as expected in the reconnecting region, may compress the gas such that the final temperature is lower than that of the ambient medium. As the entire cooling cycle occurs on a time scale during which the gas is still close to the star, the density is relatively high. Such a cooling cycle would then be similar to that proposed for R Coronae Borealis (RCB) stars (Woitke et al. 1996a). Evidently dust can be formed close to the surfaces of RCB stars, although their photospheres are much hotter than those of AGB stars. In RCB stars, dust formation is enhanced via shocks due to stellar pulsations. If flaring results in similarly strong shocks, the high densities close to the AGB surface, e.g., at radii of $\sim 1.2R_* \lesssim r \lesssim 3R_*$, could facilitate dust formation. We therefore suggest that flaring may

enhance dust formation, as long as the mass loss rate is $\dot{M} \gtrsim 10^{-6} M_{\odot} \text{ yr}^{-1}$. The condition may be somewhat different for very massive cool giants, which we do not consider here, although the same qualitative mechanism may apply.

3. OBSERVATIONAL SIGNATURES OF AGB STAR MAGNETIC ACTIVITY

As the preceding suggests, it is likely that AGB stars generate locally strong surface magnetic fields, resulting in violent reconnection events (flares). Establishing the presence of such transient magnetic activity on AGB stars observationally would have several important ramifications; in particular, we would gain new insights into stellar dynamos and the shaping of planetary nebulae. We now discuss several new observational directions that show promise for establishing the degree of magnetic activity around AGB stars.

3.1. X-ray emission and flaring

Magnetic activity — due either to differential rotation in convective regions or, for very young systems, to interactions between magnetic fields associated with star and circumstellar disk — is widely believed to be ultimately responsible for X-ray emission from late-type stars. The physical conditions that give rise to this activity, and the connection between, e.g., stellar magnetic dynamo activity and X-ray emission, remains uncertain. It is apparent, however, that highly convective late-type stars and young, low-mass stars that presumably retain high rotation velocities and/or accretion disks are among the strongest X-ray emission sources. Most low-mass, pre-main sequence (T Tauri) stars, for example, have X-ray luminosities (relative to bolometric) exceeding that of the Sun by 3-4 orders of magnitude (Feigelson & Montmerle 1999). Hence, if AGB stars indeed generate strong magnetic fields, either locally or globally, then they may be sources of X-ray emission and, perhaps, strong, long-duration (see below) X-ray flares.

3.1.1. X-ray emission from the Mira system

In addition to the apparent X-ray detection of two first-ascent red giants (Hünsch 2001), there exist two potential X-ray detections of AGB stars thus far. Both stars are in binary systems; one is a prototypical object, Mira, and the other is the symbiotic system R Aqr (Jura & Helfand 1984). The X-ray emission from R Aqr was recently imaged with the Chandra

X-ray Observatory by Kellogg et al. (2001) and appears to originate in a disk/jet system. Karovska et al. (1996) attribute the X-ray emission from the Mira system to accretion onto Mira’s companion, Mira B. Other observations also suggest the existence of an accretion disk around Mira B (e.g., Reimers & Cassatella 1985; Bochanski & Sion 2001, Wood Karovska & Raymond 2002); Jura & Helfand (1984) and Wood et al. (2002) note that it is not clear if this companion is a white dwarf or a main sequence star.

Here, we re-examine ROSAT X-ray observations⁴ of Mira (*o* Ceti), in the context of the model presented in §2. At a distance of 128 pc, Mira is one of the closest AGB stars to Earth. ROSAT detected X-ray emission toward Mira during a 9144 s pointed Position-Sensitive Proportional Counter (PSPC) observation on 1993 July 15 (Karovska et al. 1996). The background-subtracted PSPC count rate was 0.007 s^{-1} . Since only ~ 63 counts are attributed to the source (~ 10 counts are background), the archival PSPC spectrum of Mira is very noisy. Nevertheless, these data provide constraints on the nature of the X-ray source. Fitting an absorbed Raymond-Smith (RS) plasma model to the background-subtracted spectrum (Fig. 1), we find the best-fit X-ray emission temperature and absorbing column are $T_x = 1.0 \times 10^7 \text{ K}$ and $N_H = 1.8 \times 10^{21} \text{ cm}^{-2}$, respectively (with formal uncertainties of $\pm 10\%$ in T_x and $\pm 30\%$ in N_H), with a total (absorbed) model flux of $6 \times 10^{-14} \text{ ergs cm}^{-2} \text{ s}^{-1}$ (0.2-2.2 keV). The inferred intrinsic X-ray luminosity of the RS model is $\sim 2 \times 10^{29} \text{ erg s}^{-1}$, consistent with the estimate obtained by Jura & Helfand (1984). An absorbed blackbody — a model perhaps more appropriate to describe accretion-generated X-ray emission from a disk around Mira B — also produces acceptable fits; however, the temperature required for an adequate fit ($\sim 9 \times 10^5 \text{ K}$) is an order of magnitude higher than inferred via UV observations (Reimers & Cassatella 1985), and the inferred effective blackbody radius is only $\sim 10^6 \text{ cm}$. Furthermore, the value of N_H required to produce an adequate fit to the PSPC data is very large ($1.5 \pm 0.5 \times 10^{22} \text{ cm}^{-2}$). In contrast, the estimate for absorbing column obtained by fitting the RS model, $\sim 2 \times 10^{21} \text{ cm}^{-2}$, is consistent with the modest mass-loss rate of Mira as deduced from circumstellar CO radio line emission ($\sim 10^{-7} M_\odot \text{ yr}^{-1}$; e.g., Loup et al. 1993), although it is somewhat larger than the H I column density deduced from ultraviolet observations of H₂ fluorescence (Wood et al. 2002).

These results suggest that the X-ray emission is coronal in nature and, therefore, that magnetic activity is responsible for the emission. There are many possibilities as to the source(s) of this activity. It may occur on Mira A itself, or on Mira B if it is a low-mass main-sequence star (as suggested by Jura & Helfand 1984; see also Soker & Kastner 2002).

⁴These ROSAT data were obtained through the the High Energy Astrophysics Science Archive Research Center, a service of the Laboratory for High Energy Astrophysics at NASA/GSFC and the High Energy Astrophysics Division of the Smithsonian Astrophysical Observatory.

Alternatively, the emission may originate from B’s disk, either in star-disk magnetic field reconnection events or in a disk corona. Below, we further consider the implications of the first possibility, i.e., X-ray emission from Mira A.

3.1.2. *X-ray properties of AGB star flares*

If the X-ray emission detected by Einstein and ROSAT originates in magnetic flaring on Mira A, then the ROSAT results (§3.1) suggest fundamental differences between AGB star and solar flares. Using the X-ray luminosity quoted above then, assuming $n_e = 10^{10} \text{ cm}^{-3}$ and a cooling function of $\Lambda = 5 \times 10^{-23} \text{ erg cm}^3 \text{ s}^{-1}$ at $T \simeq 10^7 \text{ K}$, (Gaetz et al. 1988), we find the emitting volume to be $V_{\text{em}} = 5 \times 10^{31} \text{ cm}^3$. For a loop with scale length 10^{12} cm and cross section $(10^{11} \text{ cm})^2$, the filling factor of the emitting gas is $\epsilon \simeq 5 \times 10^{-3}$. That is, only a very small fraction of the loop is responsible for the observed X-ray emission. The filling factor can become close to unity if the electron density is lower, $n_e \sim 10^9 \text{ cm}^{-3}$. Such a low density is reasonable for Mira A, given its relatively low mass-loss rate. Even for this lower density the radiative cooling time is somewhat shorter than the expansion time, by equations (4) and (5), if the loop is smaller than the scaling in equation (5). Therefore, to hold the approximate analogy to solar flares, then the X-ray emitting flares on Mira appear to be of lower density than solar. It is assumed also that the emitting gas resides in the magnetic flux loop prior to reconnection, and there is no need for it to flow from the photosphere to a distance of $\sim 10^{12} \text{ cm}$ as the flare starts. Here, indeed, the strongly pulsating atmosphere of AGB stars is markedly different from the solar atmosphere.

In §2 we found that magnetic flares close to the the surfaces of AGB stars, and other cool giant stars, have scales of $\gtrsim 0.2R_* \sim 5 \times 10^{12} \text{ cm}$. If magnetic field reconnection proceeds at the sound speed of the gas heated to $\gtrsim 10^6 \text{ K}$, which is some fraction of the Alfven speed prior to reconnection, then the duration of a typical flare is $\tau_{\text{flare}} \gtrsim 5 \text{ days}$. A long flare might last for of order a year. This is in contrast to flares on solar-like stars, which have typical time scales of $1 - 10^4 \text{ s}$ (e.g., Pettersen 1989). Another source of flares, consisting of transient, bright emission from the optical through the X-ray regimes, can be accretion events onto close companions, including brown dwarfs and massive planets (Struck, Cohanin & Willson 2002). (The latter mechanism likely can’t account for the X-ray properties of the Mira system, however, as X-ray emission generated via accretion onto a low-mass companion would be softer than that observed.) As with magnetic flares on main sequence stars and brown dwarfs, the typical flare timescales of such events should be very short compared with those expected from magnetic flares on AGB stars. It is clear, therefore, that long-duration X-ray observations of AGB stars are very desirable as a means of establishing the presence

and timescales of flares and, hence, distinguishing between potential flare sources.

We also concluded in the last section that for high enough mass loss rate, typically $\dot{M} \gtrsim 10^{-6} M_{\odot} \text{ yr}^{-1}$, flares may enhance dust formation; in Mira, for example, the mass loss rate is too low for flare-enhanced dust formation, even if the X-ray emission originates with magnetic activity on the AGB star. The magnetic fields enhance dust formation mainly via the formation of cool spots (e.g., Soker 1998; Soker & Clayton 1999). Once local dust formation occurs, further dust formation can occur because of shadowing of the stellar radiation (Woitke, Sedlmayr, & Lopez 2000; Soker 2000), hence further increasing the inhomogeneity of the wind. Thus, in addition to searching for evidence for flaring via X-ray observations of AGB stars, it would be desirable to conduct coordinated X-ray and infrared observing campaigns designed to detect enhanced dust formation episodes during flares.

3.2. Polarized radio-line maser emission: magnetic clouds in AGB winds?

Estimates of magnetic field intensities from polarization in maser spots around AGB stars give high values (e.g., Vlemmings et al. 2002). Although the exact estimated values of the magnetic field are uncertain (Elitzur 1996), the estimates are still accurate to an order of magnitude (Elitzur, M., private communication, 2002), and in many maser spots implies that the magnetic pressure is much larger than the thermal pressure (Vlemmings et al. 2002; Bains et al. 2003). In the solar wind, on average, the magnetic pressure does not exceed the thermal pressure. However, the magnetic pressure does exceed the thermal pressure in magnetic clouds (e.g., Burlaga 2001; Yurchyshyn et al. 2001). Magnetic clouds are formed by impulsive mass loss events from the sun, and they are characterized by stronger than average magnetic fields, low proton temperatures, and smooth rotation of the magnetic field direction (e.g., Burlaga 2001). We suggest that the maser spots with strong magnetic fields which are observed around some AGB stars are similar in nature to the magnetic clouds in the solar wind, in that they represent local enhancements of the magnetic field (Soker 2002; see also Dorch & Freytag 2002 for a possible local dynamo in Betelgeuse). On average, we expect the magnetic pressure in the wind of AGB stars to be much below the thermal pressure, as winds from AGB stars are driven by radiation pressure on dust, rather than magnetic activity as in the sun.

4. SUMMARY

Recently, evidence has accumulated for the presence of relatively strong magnetic fields around AGB and similar stars. We have investigated certain theoretical and observational consequences of magnetic activity in such cool giant stars. The main results of this investigation are as follows.

1. In contrast to the solar wind, in the expanding circumstellar envelopes of AGB stars the post-flare gas cooling time is much shorter than the typical flow timescale. As a result, we predict that energetic flaring will not inhibit dust formation around AGB stars, and may even enhance dust formation in some cases.
2. Magnetic reconnection events near the stellar surface should lead to localized, long duration (timescales $\sim$ few days to a year) flares.
3. X-ray observations should provide indications as to whether AGB stars display magnetic flare activity. Analysis of an archival ROSAT X-ray spectrum of the Mira system yields results for intrinsic X-ray luminosity ($\sim 2 \times 10^{29}$ erg s $^{-1}$) and temperature ($\sim 10^7$ K) that are consistent with such activity, either on Mira A or its companion, Mira B, although we do not rule out the possibility that the X-ray emission arises from an accretion disk around Mira B. X-ray spectroscopic observations with XMM-Newton, whose sensitivity and spectral resolution far exceed those of ROSAT/PPSPC, should better constrain the X-ray emission mechanism of the Mira system. Meanwhile, the exceptional spatial resolution of the Chandra X-ray Observatory likely provides the only means to determine unambiguously the source of the X-ray emission. The separation between Mira A and B, $\sim 0.6''$ (Karovska et al. 1997), is just at the limit of Chandra’s spatial resolving power and absolute astrometry.
4. Observations of polarized maser emission from the inner circumstellar envelopes of AGB stars may indicate the presence of localized, highly magnetized wind clumps — analogous to magnetic clouds in the solar wind — rather than large-magnitude global magnetic fields.

We thank an anonymous referee for comments which led us to clarify many points. We acknowledge support for this research provided by NASA/CXO grants GO0–1067X and GO2–3009X to RIT. N.S. acknowledges support from the US-Israel Binational Science Foundation and the Israel Science Foundation.

REFERENCES

- Aschwanden, M. J., Tarbell, T. D., Nightingale, R. W., Schrijver, C. J., Title, A., Kankelborg, C. C., Martens, P., & Warren, H. P. 2000, *ApJ*, 535, 1047
- Bains, I., Gledhill, T. M., Yates, J. A., & Richards, A. M. S. 2003, *MNRAS*, in press.
- Blackman, E. G., Frank, A., Markiel, J. A., Thomas, J. H., & Van Horn, H. M. 2001, *Nature*, 409, 485
- Bochanski, J. J., & Sion, E. M. 2001, *AAS*, 198, 1208
- Burlaga, L. F. 2001, *P&SS*, 49, 1619
- Chen, P. F., Fang, C., Tang, Y. H., & Ding, M. D. 1999, *ApJ*, 513, 516
- Chevalier, R. A., & Luo, D. 1994, *ApJ*, 421, 225
- Dorch, S. B. F., & Freytag, B. 2002, in *Modeling of Stellar Atmospheres*, IAU Sym., eds., N. E. Piskunov, W. W. Weiss, and D. F. Gray, (astro-ph/0208523)
- Dorfi, E. A., & Höfner, S. 1996, *A&A*, 313, 605
- Elitzur, M. 1996, *ApJ*, 457, 415
- Elitzur, M., Ivezic, Z., & Vinkovic, D. 2002, in *Workshop on Mass-Losing Pulsating Stars and their Circumstellar Matter*, in press (astro-ph/0207513)
- Falceta-Goncalves, D., & Jatenco-Pereira, V. 2002, *ApJ*, 576, 976
- Feigelson, E. D., & Montmerle, T. 1999, *ARAA*, 37, 363
- Gaetz, T. J., Edgar, R. J., & Chevalier, R. A. 1988, *ApJ*, 329, 927
- García-Segura, G. 1997, *ApJ*, 489, L189
- García-Segura, G. 2002, in *Planetary Nebulae and their Role in the Universe*, IAU Sym. 209, ASP Conference Series, in press (astro-ph/0202041)
- García-Segura, G., Langer, N., Rozyczka, M., & Franco, J. 1999, *ApJ*, 517, 767
- García-Segura, G., & López, J. A. 2000, *ApJ*, 544, 336
- García-Segura, G., López, J. A., & Franco, J. 2001, *ApJ*, 560, 928
- Gardiner, T. A., & Frank, A. 2001, *ApJ*, 557, 250
- Hünsch, M. 2001, in *Astronomical Gesellschaft Abstract Ser.*, Vol. 18, MS 07 10
- Hünsch, M., Schmitt, J. H. M. M., Schröder, K.-P., & Zickgraf, F.-J. 1998, *A&A*, 330, 225
- Jura, M., & Helfand, D.J. 1984, *ApJ*, 287, 785
- Karovska, M., Raymond, J., & Guinan, E. 1996, Technical Report, Smithsonian Astrophysical Observatory

- Kellogg, E., Pedelty, J. A., & Lyon, R. G. 2001, *ApJ*, 563, L151
- Loup, C., Forveille, T., Omont, A., & Paul, J. F. 1993, *A&AS*, 99, 291
- Kemball, A. J., & Diamond, P. J., 1997, *ApJ*, 481, L111
- Mastrodemos, N., & Morris, M. 1999, *ApJ*, 523, 357
- Matt, S., Balick, B., Winglee, R., & Goodson, A. 2000, *ApJ*, 545, 965
- Miranda, L. F., Gomez, Y., Anglada, G., & Torrelles, J. M. 2001, *Nature*, 414, 284
- Palen, S., & Fix, J. D. 2000, *ApJ*, 531, 391
- Pascoli, G. 1997, *ApJ*, 489, 946
- Pettersen, B. R. 1989, *SoPh*, 121, 299
- Reimers, D., & Cassatella, A. 1985, *ApJ*, 297, 275
- Reimers, C., Dorfi, E. A., & Höfner, S. 2000, *A&A*, 354, 373
- Schröder, K.-P., Hünsch, M., & Schmitt, J. H. M. M. 1998, *A&A*, 335, 591
- Soker, N. 1998, *MNRAS*, 299, 1242
- Soker, N. 2000, *MNRAS*, 312, 217
- Soker, N. 2001, *ApJ*, 558, 157
- Soker, N. 2002, *MNRAS*, 336, 826
- Soker, N., & Clayton, G. C. 1999, *MNRAS*, 307, 993
- Soker, N., & Harpaz, A. 1999, *MNRAS*, 310, 1158
- Soker, N., & Kastner, J.H. 2002, *ApJ*, 570, 245
- Soker, N., & Zoabi, E. 2002, *MNRAS*, 329, 204
- Struck, C., Cohanin, B. E., & Willson, L. A. 2002, *ApJ*, 572, L83
- Szymczak, M., Cohen, R. J., & Richards, A. M. S. 1999, *MNRAS*, 304, 877
- Vesecky, J. F., Antiochos, S. K., & Underwood, J. H. 1979, *ApJ*, 233, 987
- Vlemmings, W. H. T., Diamond, P. J., & van Langevelde H. J. 2002, *A&A*, 394, 589
- Woitke, P., Goeres, A, & Sedlmayr, E. 1996a, *A&A*, 313, 217
- Woitke, P., Krüger, D., & Sedlmayr, E. 1996b, *A&A*, 311, 927
- Woitke, P., Sedlmayr, E., & Lopez, B. 2000, *A&A*, 358, 665
- Wood, B. E., Karovska, M., & Raymond, J. C. 2002, *ApJ*, 575, 1057
- Yurchyshyn, V. B., Wang, H., Goode, P. R., & Deng, Y. 2001, *ApJ*, 563, 381

Zijlstra, A. A., te Lintel Hekkert, P., Pottasch, S. R., Caswell, J. L., Ratag, M., & Habing, H. J. 1989, *A&A*, 217, 157




Fig. 1.— Archival ROSAT PSPC spectrum of the Mira system (diamonds), overlaid with best-fit absorbed Raymond-Smith plasma model (histogram). Error bars are 1σ .